



Patent claims

1. polycyclic pyrimidine-2,4(1H,3H)-diones with functionalized alkyl groups in the 1-, 3-, or both positions with the general structures Ia and Ib,



where:

R¹ is hydrogen, methyl, or ethyl

R² is hydrogen or methyl

R³ is mercapto or hydroxyaminoalkylthio (-SAIkCONHOH)

Alk is alkyl (C₁-C₅, branched and unbranched)

R₄ is hydrogen, benzyl, or phenyl

n is 0, 1 or 2,

Alk* is alkylene {C₂-C₁₂; branched or unbranched, with the exception of 3-methylpropylene [-CH₂-CH₂-CH(CH₃)-]}

X is mercapto or hydroxyaminoacylalkylthio (-SAIkCONHOH)

A is an annealed benzene ring



R⁵ is hydrogen, 6-methyl, 8-methyl, 6-fluoro, 6-chloro, 6-bromo, 6-methylthio, or 6,7-dimethoxy,

or

a 2,3-annealed thiophene ring, which may in the 4,5-positions be substituted with methyl groups or annealed with a cyclopentene, cyclohexene, or cycloheptene rings,

as well as the tautomers and pharmacologically relevant salts of these compounds.


- Sub
A2
-
2. Chinazolin-2,4(1H,3H)-diones with functionalized alkyl groups in the 1-, 3-, or both positions as described in Claim 1 for the general structures IIa and IIb



with R^1 , R^2 , R^3 , R^4 , R^5 , Alk, Alk*, n, and X defined as in Claim 1, including their tautomers and pharmacologically relevant salts.


3. The methods for the synthesis of compounds with general structures Ia, Ib, IIa, and IIb, as described in Claims 1 and 2, are distinguished by the following procedures:

(A) The reaction of 2-(alkenylamino)-1-carbonic acid amides of the general structure III,



where R^6 is alkenyl (C_3-C_6), for example allyl, methallyl, crotyl, 1-buten-4-yl, 3-penten-1-yl, and 3-hexen-1-yl, where A and R^5 are as defined above, with a $-C=S$ structural element-donating reagent, such as thiophosgene, thiourea, ammonium or alkylthiocyanate/hydrochloric acid, 1,1'-thiocarbonylbisimidazole, or preferably benzoylthiocyanate in a polar aprotic solvent,

- Sub
A2
- stir the reaction mixture
 - remove solvent under vacuum,
 - add dilute alkali solution and heat gently to ca. $60^\circ C$,
 - separate the insoluble material by filtration,
 - cool and acidify the filtrate,
 - heat the resulting compound with the general structure IV,




(IV)

where A, R_5 , and R_6 are as defined above,

- with concentrated mineral acid, such as hydrochloric acid, hydrobromic acid and/or sulfuric acid, or mixtures of these mineral acids with glacial acetic acid and or formic acid to reflux,
- cool the reaction mixture,
- dry the recovered compound of the general structure V,


Sub
A2



where

R^7, R^8, R^9, R^{10} are hydrogen, methyl, or ethyl, and A and R^5 are as defined above,

- or dry the recovered compound of the general structure VI,



(VI)


where R^{11} , R^{12} , and R^{13} are hydrogen, methyl, or ethyl, and A and R^5 are as defined above,

- in a vacuum desiccator over potassium hydroxide or
- stir the above compounds in a dilute aqueous sodium carbonate solution,
- isolate the compound,
- heat the compound with dilute mineral acid, such as hydrochloric acid,
- hydrobromic acid and or sulfuric acid, or mixtures of these mineral acids with glacial acetic acid and/or formic acid to reflux,
- cool the reaction mixture,
- wash and dry the crystallized product to obtain compounds of the general structure Ia or its tautomers, where R^4 is without exception hydrogen.

The method described in this invention for the preparation of compounds of the general structure Ia,

where R^3 is mercapto, R^4 is hydrogen, Alk, benzyl, or phenyl, $n=1$ or 2, and Alk, A, and R^5 are as defined above, is characterized by the following procedures:


Sub
Ar
B) The reaction of bi- and tricyclic 3-alkyl (or benzyl or phenyl)-pyridin-4(3H)-on-2(1H)-thiones of the general structure VII,



(VII)

where R^4 is hydrogen, benzyl, phenyl, or Alk, and A and R^5 are as described above,

with 1, ω -dihalogenalkanes of the general structure VIII,



where


m is 2, 3, or 4, and Hal is chlorine, bromine, or iodine,

- Sub
A₂
- in an aprotic dipolar solvent, preferably dimethylformamide with the addition of potassium carbonate, with extensive stirring at room temperature,
 - add dilute hydrochloric acid,
 - heat slowly and thoroughly to reflux,
 - filter the reaction mixture,
 - cool and store the filtrate at 4°C,
 - yields the compound of structure Ia,

where R³ is mercapto, R⁴ is hydrogen, Alk, benzyl, or phenyl; and Alk, A, and R⁵ are as defined above,

or

C) the reaction of 2-ammonium carboxylic acid methyl ester thiocyanate of the general structure IX,



where A and R⁵ are as defined above,

with 1,ω-dihaloalkanes of the general structure VIII,

where

m is 2 or 3, and Hal is chlorine, bromine, or iodine,

- heat the reaction partners with stirring until reflux,
- cool and isolate the precipitate,
- wash with diethylether and dry,
- dissolve the precipitate with water,
- filter the solution and add dilute sodium hydroxide to pH 10,
- isolate the precipitate and wash with water,
- dry the precipitate and shake intensively with chloroform,
- isolate the precipitate and dry,
- recrystallize from solvent, preferably 2-methoxyethanol,

to obtain compounds with the general formula V,

obtained from reaction with dibromomethane,

where R^7 , R^8 , R^9 , and R^{10} are hydrogen, and A and R^5 are as described above,

or the compound of the general structure VI,

obtained from reaction with 1,3-dibromopropane

where R^{11} , R^{12} , and R^{13} are hydrogen, and A and R^5 are as described above.

Heat the compounds obtained above of the general structures V or VI in dilute mineral acids, such as hydrochloric acid, hydrobromic acid and or sulfuric acid, or mixtures of these mineral acids with glacial acetic acid and/or formic acid to reflux,

- cool the reaction mixture,
- wash and dry the crystallized material to obtain compounds of the general structure Ia or its tautomers,


where R^1 , R^2 , R^4 are hydrogen, R^3 is mercapto, $n = 1$ or 2 , and A and R^5 are as defined above.

The method described in this invention for the preparation of compounds of the general structure Ib,

where Alk* is n-butylene ($-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$), X is mercapto, and A and R^5 are as defined above,

is characterized by the following procedures:


D) The reaction of 2-isothiocyanato-1-carboxylic acid ester of the general structure X,



(X)

where Alk, A, and R^5 are as defined above,
with 4 aminobutan-1-ol at room temperature with intensive and extensive stirring


- add water to obtain compounds of the general structure XI,



(XI)

where A and R⁵ are as defined above,

- react compounds of the general structure XI with concentrated mineral acids, such as hydrochloric acid, hydrobromic acid and or sulfuric acid, or mixtures of these mineral acids with glacial acetic acid and/or formic acid to reflux,
- cool the reaction mixture and isolate the crystalline precipitate,
- add aqueous sodium carbonate to the precipitate to adjust pH to ca. 9,
- isolate the crystalline precipitate, wash with water, and dry to obtain polycyclic 2,3,4,5-tetrahydro-7H-[1,3]thiazepin[2,3-a]pyrimidin-7-one or the general structure XII,



(XII)

where A and R⁵ are as defined above,

Heat the above mentioned compounds of the general structure XII in extremely dilute mineral acids, such as hydrochloric acid, hydrobromic acid and or sulfuric acid, or mixtures of these mineral acids with glacial acetic acid and/or formic acid to reflux,

- cool the reaction mixture,
- wash and dry the crystalline precipitate


to obtain compounds of the general structure Ib or their tautomers,

where Alk* is n-butylene (-CH₂-CH₂-CH₂-CH₂-), X is mercapto, and A and R⁵ are as defined above.

The method described in this invention for the preparation of compounds of the general structure Ib, where Alk* is alkylene {C₂-C₁₂; branched or unbranched, with the exception of 3-methylpropylene [-CH₂-CH₂-CH(CH₃)-

]} , X is mercapto, and A and R⁵ are as defined above, is characterized by the following procedures:

E) The reaction of 2-alkoxycarbonylamino-1-carboxylic acid alkyl ester of the general structure XIII,



where Alk is alkyl (C₁-C₃) and A and R⁵ are as defined above.


with aminoalkanols of the general structure XIV,

where Alk* is alkylene {C₂-C₁₂; branched or unbranched, with the exception of 3-methylpropylene [-CH₂-CH₂-CH(CH₃)-] },

by heating in well-known reactions,

- cool the reaction mixture,
- add water and dilute hydrochloric acid to ca. pH 4,

- isolate the precipitate, wash with water, and dry to obtain the polycyclic 3-(ω -hydroxyalkyl)-pyrimidine-2,4(1H,3H)-diones of the general structure XV,




(XVI)

where Alk* is alkylene {C₂-C₁₂; branched or unbranched, with the exception of 3-methylpropylene [-CH₂-CH₂-CH(CH₃)-] } and A and R⁵ are as defined above.

React the compounds of the general structure XV with concentrated hydrochloric acid, phosphorus trichloride, or phosphoroxidchloride, but preferably with concentrated hydrobromic acid by

- heating to reflux,
- cooling the reaction mixture,

- isolating the precipitate, washing with water and drying to obtain polycyclic 3- (ω-haloalkyl)-pyrimidin-2,4(1H,3H)-diones of the general structure XVI,



(XVII)

where Hal is chlorine, or bromine, and Alk*, and A and R⁵ are as defined above.

React the compounds of the general structure XVI in rapidly boiling polar solvent with thiourea,

- heat to reflux,
- cool the reaction mixture,
- add water and dilute aqueous sodium hydroxide to obtain an alkaline pH,
- after the solution clears, filter it and
- add to the filtrate dilute hydrochloric acid to ca. pH 3,

- isolate the precipitate, wash with water, and dry to obtain compounds of the general structure Ib or their tautomers

where Alk* is alkylene (C₄-C₁₂; branched or unbranched, with the exception of 3-methylpropylene [-CH₂-CH₂-CH(CH₃)-] }, X is mercapto, and A and R⁵ are as defined above.

Sub
A²
The method described in this invention for the preparation of compounds of the general structure Ia or Ib, where R³ or X is hydroxaminoacylalkylthio (-SAIkCONHOH) and R¹, R², R⁴, Alk, n, Alk*, A, and R⁵ are as defined above,

is characterized by the following procedures:

F) React compounds of the general structure Ia or Ib,

where R³ is mercapto, and R¹, R², R⁴, Alk, n, Alk*, A, and R⁵ are as defined above,

with N-hydroxyhalogencarboxylic acid amide, preferably 2-chloro-N-hydroxyacetamide (ClCH₂CONHOH),

- either in pyridine
- or in acetone solution, preferable in the presence of base,
- at room temperature,

- remove solvent by distillation,
- add water,
- isolate the precipitate, wash with water and dry to obtain compounds with the general structure Ia or Ib or their tautomers,

where R^3 or X is hydroxyaminoacylalkylthio,

500
A2
4. methods based on claim 3, characterized by one or more of the following steps:

- a) the reaction of compounds of general structure III with benzoylisothiocyanate in absolute acetone.
- b) the reaction of compounds of general structure VII with compounds of general structure VIII in absolute dimethylformamide with dry potassium carbonate.

5. Pharmaceutical preparations containing one or more compounds in claim 1 and or 2, as well as commonly used filler and inert ingredients, and binders.

6. Pharmaceutical preparations with collagenase/MMP-inhibiting activities,

Sub

thus characterized by the fact that they contain as active ingredients

A2

one or more compounds covered by claims 1 and 2.
